

Law of large numbers for superdiffusions: the non-ergodic case

János Engländer

Department of Statistics and Applied Probability
University of California, Santa Barbara
CA 93106-3110, USA.

Email: `englander@pstat.ucsb.edu`

February 2, 2008

Abstract

In [8] the Law of Large Numbers for the local mass of certain superdiffusions was proved under a spectral theoretical assumption, which is equivalent to the ergodicity (positive recurrence) of the motion component of an H -transformed (or weighted) superprocess. In fact the assumption is also equivalent to the property that the scaling for the expectation of the local mass is pure exponential.

In this paper we go beyond ergodicity, that is we consider cases when the scaling is not purely exponential. *Inter alia*, we prove the analog of the Watanabe-Biggins Law of Large Numbers for super-Brownian motion (SBM).

We will also prove another Law of Large Numbers for a bounded set moving with subcritical speed, provided the variance term decays sufficiently fast.

Further illustrative examples, such as SBM with drift and super-Ornstein-Uhlenbeck process, will be provided too.

MSC 2000 subject classifications. 60J60, 60J80

Key words and phrases. super-Brownian motion, super-Ornstein-Uhlenbeck process, superdiffusion, superprocess, Law of Large Numbers, H -transform, weighted superprocess, scaling limit, local extinction, local survival.

1 Introduction and statement of results

1.1 Basic notions

Let $D \subseteq \mathbb{R}^d$ be a domain and let $\mathcal{B}(D)$ denote the Borel sets of D . We write $\mathcal{M}_f(D)$ and $\mathcal{M}_c(D)$ for the class of finite measures resp. the class of finite measures with compact support on $\mathcal{B}(D)$. For $\mu \in \mathcal{M}_f(D)$, denote $\|\mu\| := \mu(D)$ and let $C_b^+(D)$ and $C_c^+(D)$ be the class of non-negative bounded continuous resp. non-negative continuous functions $D \rightarrow \mathbb{R}$ having compact support. Write $C^{k,\eta}(D)$ for the usual Hölder spaces of index $\eta \in (0, 1]$ including derivatives of order k , and set $C^\eta(D) := C^{0,\eta}(D)$. Let L be an elliptic operator on the domain $D \subseteq \mathbb{R}^d$ of the form

$$L := \frac{1}{2} \nabla \cdot a \nabla + b \cdot \nabla, \quad (1)$$

where $a_{i,j}, b_i \in C^{1,\eta}(D)$, $i, j = 1, \dots, d$, for some $\eta \in (0, 1]$, and the matrix $a(x) := (a_{i,j}(x))$ is symmetric, and positive definite for all $x \in D$. In addition, let $\alpha, \beta \in C^\eta(D)$, and assume that α is positive, and β is bounded from above.

Definition 1 ($(L, \beta, \alpha; D)$ -superdiffusion) With D, L, β and α as above, let $(X, \mathbf{P}^\mu, \mu \in \mathcal{M}_f(D))$ denote the $(L, \beta, \alpha; D)$ -superdiffusion. That is, X is the unique $\mathcal{M}_f(D)$ -valued continuous (time-homogeneous) Markov process which satisfies, for any $g \in C_b^+(D)$,

$$\mathbf{E}^\mu \exp \langle X_t, -g \rangle = \exp \langle \mu, -u(\cdot, t) \rangle, \quad (2)$$

where u is the minimal nonnegative solution to

$$\left. \begin{aligned} u_t &= Lu + \beta u - \alpha u^2 \quad \text{on } D \times (0, \infty), \\ \lim_{t \downarrow 0} u(\cdot, t) &= g(\cdot). \end{aligned} \right\} \quad (3)$$

As usual, $\langle \nu, f \rangle$ denotes the integral $\int_D \nu(dx) f(x)$.

(See Dynkin (1991, 2002) [3], [4] or Dawson (1993) [2] for the definition of superprocesses in general; see Engländer and Pinsky (1999) [5] for more on the definition in the particular setting above.)

One usually refers to β as *mass creation* and α as the *intensity parameter* (or variance).

Note that although the above model is a time-homogeneous process, later on the introduction of certain *time-inhomogeneous* superdiffusions will be necessary.

1.2 Law of Large Numbers vs. Local extinction

Our principal interest is in establishing a Law of Large Numbers for the local mass of certain superdiffusions. That is if $f \in C_c^+(D) \not\equiv 0$ and $\mu \in \mathcal{M}_c(D) \not\equiv 0$, then we would like to verify that,

$$\lim_{t \rightarrow \infty} \frac{\langle X_t, f \rangle}{\mathbf{E}^\mu \langle X_t, f \rangle} = N_\mu,$$

where N_μ is a (non-degenerate) random variable and the limit holds in some suitable sense. (See [8] for an explanation on why such a statement can be called a Law of Large Numbers.)

One can immediately see however, that such a law cannot be true without having some condition on the operator corresponding to the superprocess. To elucidate on this point, note that the Law of Large Numbers will obviously not hold when X *exhibits local extinction* (i.e., the support of X leaves any given bounded set, $\mathbf{P}^\mu$ -a.s. for each $\mu \in \mathcal{M}_c$). Now, let

$$\lambda_c = \lambda_c(L + \beta) = \inf\{\lambda \in \mathbb{R} : \exists u > 0 \text{ satisfying } (L + \beta - \lambda)u = 0 \text{ in } D\} \quad (4)$$

denote the *generalized principal eigenvalue* for $L + \beta$ on D (see the Appendix). Since Pinsky proved that X exhibits local extinction if and only if $\lambda_c \leq 0$, we can only hope to have the Law of Large Numbers if we assume that $\lambda_c > 0$.

1.3 Motivation

In order to understand what follows, we need to present first some concepts regarding the *criticality theory of second order elliptic operators* (for a complete presentation, the reader is referred to Chapter 4 in [10]). The operator $L + \beta - \lambda_c$ is called *critical* if there exists a positive function f satisfying that $(L + \beta - \lambda_c)f = 0$ but there is no (minimal positive) Green's function for the operator $L + \beta - \lambda_c$. In this case f is unique up to constant multiples and is called the *ground state*. The operator $L + \beta - \lambda_c$ is called *product-critical* if it is critical with ground state $0 < \phi_c$, and ϕ_c and $\tilde{\phi}_c$ (i.e. the ground state for the formal adjoint of $L + \beta - \lambda_c$) satisfy $\langle dx, \phi_c \tilde{\phi}_c \rangle < \infty$. In this case we normalize them by $\langle dx, \phi_c \tilde{\phi}_c \rangle = 1$. If $L + \beta - \lambda_c$ possesses a Green's function, then it is called *subcritical*.

The following has been shown in [8]:

Proposition 1 ([8], Theorem 1) *In addition to the assumption $\lambda_c > 0$, also assume that $L + \beta - \lambda_c$ is product-critical, that $\alpha\phi_c$ is bounded and that*

X starts in a state μ with $\langle \mu, \phi_c \rangle < \infty$. Let $f \in C_c^+(D)$. If $f \not\equiv 0$ and $\|\mu\| \neq 0$, then

$$\lim_{t \rightarrow \infty} \frac{\langle X_t, f \rangle}{\mathbf{E}^\mu \langle X_t, f \rangle} = \frac{\overline{X}_\infty^H}{\langle \mu, \phi_c \rangle}, \quad \text{in } \mathbf{P}^\mu\text{-probability.} \quad (5)$$

A simple case of a superdiffusion is when $D = \mathbb{R}^d$, $d \geq 1$, $L = \frac{1}{2}\Delta$, with α, β positive constants (supercritical super-Brownian motion). Here $\lambda_c = \beta$ and

$$\frac{1}{2}\Delta + \beta - \lambda_c = \frac{1}{2}\Delta.$$

Since $\phi_c = \tilde{\phi}_c \equiv 1$, $d \geq 1$, the operator $\frac{1}{2}\Delta$ is either critical but not product-critical ($d \leq 2$), or subcritical ($d \geq 3$). Therefore *this case was not included in the setup* of [8]. On the other hand, the corresponding (Strong) Law of large Numbers is well known for *discrete particle systems*.

Using techniques from Fourier transform theory, Watanabe [12] proved SLLN for branching-Brownian motion in $\mathbb{R}^d$ and in certain subdomains of it. It is not clear however if his method can be generalized for more general branching diffusions. Furthermore, the proof in [12] is thought to have a gap. In [12] a family of nonnegative martingales $\{W_t^\lambda; \lambda \in \Lambda\}$ is considered, together with the family of their limits $\mathfrak{L} := \{W^\lambda; \lambda \in \Lambda\}$, where $W^\lambda := \lim_{t \rightarrow \infty} W_t^\lambda$. The problem, however, is that those limits are only *almost sure* ones, so for every $\lambda \in \Lambda$ there is an exceptional null set N_λ . Since Λ is not countable, $\mathfrak{L}$ is not defined on the *uncountable* union $\bigcup_{\lambda \in \Lambda} N_\lambda$!

Biggins [1] gives a complete proof for the case when branching-Brownian motion is replaced by branching RW. Here time is discrete: $n = 1, 2, \dots$ and instead of considering $\{W_n^\lambda\}$ as a family indexed by λ , it is thought of as a sequence of (continuous) functions $\{W_n(\cdot)\}$, i.e. λ is not the index but the argument. If the sequence is restricted to a compact subset $F \subset \Lambda$, it can be thought of as a martingale with values in the Banach space of continuous functions on F (under the supremum norm). Biggins then proves that this martingale converges almost surely and in mean.

The purpose of this paper is to prove the Law of Large Numbers for a class of superprocesses that *includes supercritical super-Brownian motion*. Instead of trying to adapt the Watanabe-Biggins approach to our setting, our method will use some ingredients from [8], however the time scales will have to be modified now and also the spatial spread of the process must be handled. In particular, to verify that supercritical super-Brownian motion is included in the setup (satisfies the assumption on the spatial speed), we will need a result from [11] too.

1.4 Main results

Throughout the paper the following assumption will be in force.

Assumption 1 Let $\lambda_c = \lambda_c(L + \beta; D)$ denote the generalized principal eigenvalue of $L + \beta$ on D , and let $S = \{S_t\}_{t \geq 0}$ denote the semigroup corresponding to $L + \beta - \lambda_c$ on D .

(A.1) **(local survival)** $\lambda_c > 0$,

(A.2) **(scaling of linear semigroup)** There exist two functions $s : (0, \infty) \rightarrow (0, \infty)$, and $h : D \rightarrow (0, \infty)$, $h \in C^{2,\eta}(D)$, and a locally finite measure $r(dx)$ such that

- (a) $\log s_t = \mathcal{O}(\log t)$ (i.e. $\sup_{t>0} \frac{\log s_t}{\log t} < \infty$) and $\lim_{t \rightarrow \infty} (\log s_t)' = 0$,
- (b) $\sup_D \alpha h < \infty$,
- (c) $\lim_{t \rightarrow \infty} \langle s_t \cdot S_t(f)(x), \mu(dx) \rangle = \langle f, r(dy) \rangle \cdot \langle h, \mu \rangle$ for all $f \in C_c^+(D)$, and $\mu \in \frac{1}{h} \mathcal{M}_f$,

(A.3) **(spatial spread)** There exist two more functions, $z, \widehat{z} : (0, \infty) \rightarrow (0, \infty)$ such that

- (a) $\lim_{t \rightarrow \infty} \widehat{z}_t = \infty$,
- (b) $\log(t + \widehat{z}_t) = o(t)$, $t \rightarrow \infty$,
- (c) $\lim_{t \rightarrow \infty} s_{t+\widehat{z}_t} / s_{\widehat{z}_t} = 1$,
- (d) $\lim_{t \rightarrow \infty} P^\mu(\langle X_t, B_{z_t}^c(0) \rangle > 0) = 0$,
- (e) if $f \in C_c^+(D)$, then

$$\lim_{t \rightarrow \infty} \sup_{|x| \leq z_t} \left| \frac{s_{\widehat{z}_t}}{h(x)} \cdot S_{\widehat{z}_t}(f)(x) - \langle f, r(dy) \rangle \right| = 0.$$

Let $H(x, t) := \exp(-\lambda_c t)h(x)$ and consider the weighted superprocess X^H (see section 3). Abbreviate $W := X^H$ and let $\overline{W}$ denote the *total mass process*, i.e.

$$\overline{W} = \overline{X}^H := \|X^H\|. \quad (6)$$

It is not hard to show that $\overline{W}$ is a supermartingale (see the Appendix), and therefore it has a limit, $\overline{W}_\infty$. One does not have $\mathbf{E}^\mu \overline{W}_\infty = \langle \mu, h \rangle$ in general. However, when L_0^h is conservative on D , that is, it never leaves the domain D with probability one, $\overline{W}$ is a UI martingale (see again the Appendix). In this case, by uniform integrability, $\mathbf{E}^\mu \overline{W}_\infty = \langle \mu, h \rangle > 0$.

Theorem 1 (Law of Large Numbers) *With the notations of Assumption 1, if $f \in C_c^+(D) \not\equiv 0$ and $\mu \in \mathcal{M}_c(D) \not\equiv 0$, then in probability,*

$$\lim_{t \rightarrow \infty} \frac{\langle X_t, f \rangle}{\mathbf{E}^\mu \langle X_t, f \rangle} = \frac{\overline{W}_\infty}{\langle \mu, h \rangle}.$$

The limit is mean-one (and in particular, not identically zero) when L_0^h corresponds to a conservative diffusion.

In order to give a simple condition for the limit to be mean-one, we recall the *compact support property*.

Definition 2 *The $(L, \beta, \alpha; D)$ -superdiffusion possesses the compact support property if*

$$\mathbf{P}^\mu \left(\bigcup_{0 \leq s \leq t} \text{supp } X_s \subset\subset D \right) = 1, \text{ for all } \mu \in \mathcal{M}_c(D), t \geq 0. \quad (7)$$

[Here $A \subset\subset D$ means that the closure of the bounded domain A is in D .] Since there are various conditions given in [5, 6] for the compact support property to hold, the following result is useful.

Theorem 2 (No loss of mass in the limit) *If the compact support property holds, then the diffusion process corresponding to L_0^h on D is conservative, and consequently, the limit appearing in Theorem 1 is mean-one.*

Using H -transforms we will also prove another Law of Large Numbers for a bounded set moving with subcritical speed, provided the variance term α decays sufficiently fast.

Theorem 3 (Law of Large Numbers for a moving bounded set) *Let $c < \sqrt{2\beta}$, $\alpha(x) \leq K \exp(-cx)$ with some $K > 0$, and define $f^{(ct)}(x) := f(x_1 + ct, x_2, \dots, x_d)$. With the notations of Assumption 1, if $f \in C_c^+(D) \not\equiv 0$ and $\mu \in \mathcal{M}_c(D) \not\equiv 0$, then in probability,*

$$\lim_{t \rightarrow \infty} \frac{\langle X_t, f^{(ct)} \rangle}{\mathbf{E}^\mu \langle X_t, f^{(ct)} \rangle} = N_c,$$

where N_c is a nonnegative, mean one random variable.

1.5 Outline

The rest of this paper is organized as follows. In section 2 we give examples for our main theorem. In section 3 we give a review on a fundamental tool, a space-time transformation. In section 4 we present the proofs. Finally, the Appendix summarizes the necessary background.

2 Examples

In this section we give five examples. All of them satisfy the assumption, and thus obey the Law of Large Numbers. In all the examples, $D = \mathbb{R}^d$, and the constant c (appearing in all but the first example) is positive. In our notation x is the d -dimensional vector $(x_1, x_2, \dots, x_d)$ and $x^2 := |x|^2 = \sum_{i=1}^d x_i^2$. All the examples are versions of either the super-Brownian motion (SBM) or the super-Ornstein-Uhlenbeck process (SOU).

Example 1 (supercritical SBM) The assumptions are satisfied for supercritical super-Brownian motion. Indeed, if $\beta(\cdot) \equiv \beta > 0$, then $\lambda_c = \beta$, because $\lambda_c(\Delta, \mathbb{R}^d) = 0$. Furthermore choose $h \equiv 1$, $r(dx) := dx$ and the Brownian scaling factor $s_t := t^{d/2}$. Finally, as far as the spatial spread of the process is concerned, z_t can be defined as $z_t := (\sqrt{2\beta} + \epsilon)t$ (see [11]), and thus $\hat{z}_t$ can be defined e.g. as $\hat{z}_t = t^m$ with $m > 2$. This setting satisfies conditions (2)-(3), as long as $0 < \alpha$ is bounded from above. $\diamond$

Consider now the simplest case of the previous example, the one when α is a positive constant. Then the non-degenerate random variable $\overline{W}_\infty$ can be thought of as the scaled limit of a one dimensional diffusion. Indeed, $Y := \|X\|$ is a diffusion corresponding to the operator

$$\mathcal{L} := x \left(\alpha \frac{\partial^2}{\partial x^2} + \beta \frac{\partial}{\partial x} \right) \text{ on } [0, \infty)$$

with $Y_0 = \|\mu\|$, and $\overline{W}_\infty = \lim_{t \rightarrow \infty} e^{-\beta t} Y_t$.

For the reader unfamiliar with *spatial h-transform* it is helpful to review section 3 before reading the rest of the examples. (The h -transform is a particular case of the H -transform with $H(t, x) = h(x)$.)

In the following examples there is no need to check our assumptions for the Law of Large Numbers to hold. The validity of the Law of Large Numbers will simply follow from its invariance under h -transforms and from the Law of Large Numbers in the previous example. This will in particular mean that the limiting random variable is always *non-degenerate*.

Let X be the supercritical super-Brownian motion of the first example. Since the Law of Large Numbers holds true for X starting with any nonzero finite measure, therefore it is also true for X^h starting with any measure of the form $\nu = h\mu$, where $0 \neq \mu$ is a finite measure. To avoid working with these h -dependent spaces, we will simply assume in all the examples below that the initial (nonzero) measure belongs to $\mathcal{M}_c$.

Example 2 (supercritical SBM with drift) Let $h(x) := e^{cx_1}$. Then X^h corresponds to the motion generator

$$\mathcal{L} := \frac{1}{2}\Delta + b \cdot \nabla, \quad b := (c, 0, \dots, 0)$$

(Brownian motion with drift in the first coordinate direction), mass creation $\beta^h = \beta + c^2/2 > c^2/2$ and intensity parameter $\alpha^h = \alpha e^{cx_1}$. Hence, *the super-Brownian motion with drift $c > 0$ obeys the Law of Large Numbers, if the mass creation is larger than $c^2/2$, and the intensity parameter is $\mathcal{O}(e^{cx_1})$, $|x_1| \rightarrow \infty$.* $\diamond$

Example 3 (supercritical SBM with outward drift) Let $0 < h \in C^{2,\eta}$ satisfy $h(x) := e^{c|x|}$, $|x| \gg 1$. Then X^h corresponds to the motion generator

$$\mathcal{L} := \frac{1}{2}\Delta + b \cdot \nabla, \quad b(x) = c \frac{x}{|x|}, \text{ for } |x| \gg 1,$$

mass creation $\beta^h = \beta + c^2/2 > c^2/2$ and $\alpha^h = \alpha e^{c|x|}$. Hence, *any superprocess obeys the Law of Large Numbers, if the motion component is Brownian motion with outward drift $c > 0$ and the mass creation is larger than $c^2/2$, while the intensity parameter is $\mathcal{O}(e^{c|x|})$, $|x| \rightarrow \infty$.* $\diamond$

The last two examples concern ‘super-Ornstein-Uhlenbeck processes’.

Example 4 (supercritical SOU with small α) We now choose $h(x) := e^{-cx^2}$. Then X^h corresponds to the Ornstein-Uhlenbeck migration with generator

$$\mathcal{L} := \frac{1}{2}\Delta - 2cx \cdot \nabla,$$

mass creation $\beta(x) = K + 2c^2x^2$, $K > -c$, and $\alpha^h := \alpha e^{-cx^2}$. Hence, *the super-Ornstein-Uhlenbeck process with drift $c > 0$ obeys the Law of Large Numbers, as long as the mass creation is of the above form, and the intensity parameter is $\mathcal{O}(e^{-cx^2})$, $|x| \rightarrow \infty$.* $\diamond$

Example 5 (supercritical outward SOU with large β) Taking $h(x) := e^{cx^2}$, X^h corresponds to the migration with generator

$$\mathcal{L} := \frac{1}{2}\Delta + 2cx \cdot \nabla,$$

mass creation $\beta(x) = K + 2c^2x^2$, $K > c$, and $\alpha^h = \alpha e^{cx^2}$. Hence, *any superprocess obeys the Law of Large Numbers, if the motion component is an outward drifting Ornstein-Uhlenbeck process with drift $c > 0$ and the mass creation is of the above form, while the intensity parameter is $\mathcal{O}(e^{cx^2})$, $|x| \rightarrow \infty$.* $\diamond$

We note that if in the last two example one replaces $\beta(\cdot)$ by a positive constant β , then the models will belong to the setup treated in [8] (product-critical, or ergodic case).

3 The H -transform of superdiffusions

This section gives a review on the H -transform¹. The H -transform, introduced in [8], is a space-time generalization of the spatial h -transform for superdiffusions (h -transformed, or weighted superprocesses had been introduced earlier in [5]).

We first review the more general definition of a time-inhomogeneous superdiffusion. Let $\tilde{L}$ be an elliptic operator on $D \times \mathbb{R}^+$ of the form

$$\tilde{L} := \frac{1}{2}\nabla \cdot \tilde{a}\nabla + \tilde{b} \cdot \nabla \quad (8)$$

where the functions $\tilde{a}_{i,j}, \tilde{b}_i : D \times \mathbb{R}^+ \rightarrow \mathbb{R}$, $i, j = 1, \dots, d$ are $C^{1,\eta}(D)$ (for some $\eta \in (0, 1]$) in the space, and continuously differentiable in the time coordinate. Moreover assume that the symmetric matrix $\tilde{a}(x, t) := (a_{i,j}(x, t))$ is positive definite for all $x \in D$ and $t \in \mathbb{R}^+$.

In addition, let $\tilde{\alpha}, \tilde{\beta} : D \times \mathbb{R}^+ \rightarrow \mathbb{R}$, be $C^\eta(D)$ in the space, and continuously differentiable in the time coordinate. Finally assume that $\tilde{\alpha}$ is positive, and $\tilde{\beta}$ is bounded from above.

Definition 3 (Time-inhomogeneous $(\tilde{L}, \tilde{\beta}, \tilde{\alpha}; D)$ -superdiffusion)

¹The reader should not confuse with the space-time harmonic transformation yielding a Girsanov-type change of measure – see e.g. [9].

(i) The $(\tilde{L}, \tilde{\beta}, \tilde{\alpha}; D)$ -superdiffusion is a measure-valued (inhomogeneous) Markov process, $(X, \mathbf{P}^{\mu, r}; \mu \in \mathcal{M}_f(D), r \geq 0)$, that is, a family $\{\mathbf{P}^{\mu, r}\}$ of probability measures where $\mathbf{P}^{\mu, r}$ is a probability on $C([r, \infty))$ and the family is indexed by $\mathcal{M}_f(D) \times [0, \infty)$, such that the following holds: for each $g \in C_b^+(D)$ and $\mu \in \mathcal{M}_f(D)$,

$$\mathbf{E}^{\mu, r}[\exp -\langle X_t, g \rangle] = \exp -\langle \mu, u(\cdot, r; t, g) \rangle, \quad (9)$$

where $u = u(\cdot, \cdot; t, g)$ is a particular non-negative solution to the backward equation

$$\begin{aligned} -\partial_r u &= \tilde{L}u + \tilde{\beta}u - \tilde{\alpha}u^2 && \text{in } D \times (0, t), \\ \lim_{r \uparrow t} u(\cdot, r; t, g) &= g(\cdot). \end{aligned} \quad (10)$$

(ii) To determine the solution u uniquely, use the equivalent *forward* equation along with the minimality of the solution: fix $t > 0$ and introduce the ‘time-reversed’ operator $\hat{L}$ on $D \times (0, t)$ by

$$\hat{L} := \frac{1}{2} \nabla \cdot \hat{a} \nabla + \hat{b} \cdot \nabla, \quad (11)$$

where, for $r \in [0, t]$,

$$\hat{a}(\cdot, r) := \tilde{a}(\cdot, t - r) \text{ and } \hat{b}(\cdot, r) := \tilde{b}(\cdot, t - r);$$

furthermore let

$$\hat{\beta}(\cdot, r) := \tilde{\beta}(\cdot, t - r) \text{ and } \hat{\alpha}(\cdot, r) := \tilde{\alpha}(\cdot, t - r).$$

Consider now v , the *minimal* non-negative solution to the *forward* equation

$$\begin{aligned} \partial_r v &= \hat{L}v + \hat{\beta}v - \hat{\alpha}v^2 && \text{in } D \times (0, t), \\ \lim_{r \downarrow 0} v(\cdot, r; t, g) &= g(\cdot). \end{aligned} \quad (12)$$

Then

$$u(\cdot, r; t, g) = v(\cdot, t - r; t, g).$$

(See also [8] concerning the construction of minimal non-negative solutions for forward equations.)

As we will see in Lemma 1 (b), one way of defining a time-inhomogeneous superdiffusion is to start with a time-homogeneous one, and then to apply an ‘ H -transform’. In general, the H -transform of a time-inhomogeneous superdiffusion is defined as follows. Let $0 < H \in C^{2,\eta}(D) \times C^{1,\eta}(\mathbb{R}^+)$ and let X be a $(\tilde{L}, \tilde{\beta}, \tilde{\alpha}; D)$ -superdiffusion. We define a new process X^H by

$$X_t^H := H(\cdot, t)X_t \quad \left(\text{that is, } \frac{dX_t^H}{dX_t} = H(\cdot, t) \right), \quad t \geq 0. \quad (13)$$

This way one obtains a new superdiffusion, which, in general, *is not finite measure-valued* but only σ -finite measure-valued. That is, if $\mathcal{M}(D)$ denotes the family of all (finite or infinite) measures on D , then

$$X_t^H \in \mathcal{M}_H^{(t)}(D) := \{\nu \in \mathcal{M}(D) \mid H(\cdot, t)^{-1}\nu \in \mathcal{M}_f(D)\}$$

(c.f. [5], p. 688.)

In [5], Section 2, it was shown, that, from an analytical point of view, the (spatial) h -transform of the superdiffusion is given by a certain transformation of the corresponding semilinear operator. This remains the case for the space-time H -transform. The following result was proved in [8].

Lemma 1

Let X^H be defined by (13). Then

- (a) X^H is a $(\tilde{L} + \tilde{a}\frac{\nabla H}{H} \cdot \nabla, \tilde{\beta} + \frac{\tilde{L}H}{H} + \frac{\partial_r H}{H}, \tilde{\alpha}H; D)$ -superdiffusion.
- (b) In particular, if X is a time-homogeneous $(L, \beta, \alpha; D)$ -superdiffusion, and H is of the form

$$H(x, t) := e^{-\lambda_c t} h(x), \quad (14)$$

where λ_c is the principal eigenvalue of $L + \beta$, and h is a positive solution of $(L + \beta)h = \lambda_c h$, then X^H is a $(L + a\frac{\nabla h}{h} \cdot \nabla, 0, \alpha h e^{-\lambda_c t}; D)$ -superdiffusion.

Remark 1 (Unbounded $\tilde{\beta}$'s) As it is already the case with the spatial h -transform for superdiffusions, it is possible that the coefficient $\tilde{\beta}$ transforms into a new coefficient that is no longer bounded. In fact this can be the very definition of superdiffusions for certain unbounded $\tilde{\beta}$'s (see [5], Section 2 for explanation). $\diamond$

Remark 2 (Invariance of LLN) When the function H is of the form $H(x, t) = h(x)r(t)$, the validity of the Law of Large Numbers is *invariant* under the transformation. Indeed, if f belongs to $C_c^+(D)$, then so does $g := hf$; and for $\nu = r(0)h\mu$,

$$\frac{\langle X_t^H, f \rangle}{\widetilde{\mathbf{E}}^\nu \langle X_t^H, f \rangle} = \frac{\langle X_t, g \rangle}{\mathbf{E}^\mu \langle X_t, g \rangle},$$

and the invariance follows by letting $t \rightarrow \infty$. $\diamond$

4 Proofs

4.1 Proof of Theorem 1

Having the H -transform at hand we now reformulate our assumptions and also the statement in terms of the space-time weighted superprocess X^H , where H is of the form

$$H(x, t) := e^{-\lambda_c t} h(x), \quad (15)$$

λ_c is the principal eigenvalue of $L + \beta$, and h is the function appearing in the assumption. Abbreviate

$$L_0^h := L + a \frac{\nabla h}{h} \cdot \nabla \quad (16)$$

and note that in fact

$$L_0^h(u) = h^{-1}(L + \beta - \lambda_c)(hu) = H^{-1}(L + \beta + \partial_t)(Hu).$$

Let $\mathfrak{S} = \{\mathfrak{S}_s\}_{s \geq 0}$ denote the semigroup corresponding to the operator $L_0^h = (L + \beta - \lambda_c)^h$, that is, $\mathfrak{S} := S^h$. Since the semigroup $\mathfrak{S}$ corresponds to an operator with no zeroth order part (i.e. to L_0^h),

$$\mathfrak{S}_s 1 \leq 1. \quad (17)$$

If the diffusion process corresponding to L_0^h on D is conservative, then in fact $\mathfrak{S}_s 1 = 1$; in general one only has (17). (See more on conservativeness and its connection to a uniformly integrable martingale in the Appendix).

Let us use the shorthand $u' := \partial_t u$. We claim that h is a positive solution of $(L + \beta)h = \lambda_c h$. To see this note that (A.2)(c) with $\mu = \delta_x$ yields $\lim_{t \rightarrow \infty} s_t \cdot S_t(f)(x) = \langle f, r(dy) \rangle \cdot h(x)$ for $x \in D$ and $f \in C_c^+(D)$. Then,

defining $\widehat{\beta}(t, x) := \beta(x) + (\log s_t)'$, the equation $(L + \beta)h = \lambda_c h$ follows from the fact that $u(t, x) := S_t(f)(x)$ solves

$$(L + \beta - \lambda_c)u = u'$$

and therefore $v(t, x) := s_t \cdot u(t, x)$ solves

$$(L + \beta - \lambda_c)v = v' - (\log s_t)'v,$$

that is

$$(L + \widehat{\beta} - \lambda_c)v = v'.$$

Indeed, first note that

$$\lim_{t \rightarrow \infty} \sup_{x \in D} |\widehat{\beta}(t, x) - \beta(x)| = 0.$$

Then a standard argument (see [5], p.708) together with the second relation in (A.2)(a) gives that $w(x) := \lim_{t \rightarrow \infty} v(t, x) = \langle f, r(dy) \rangle \cdot h(x)$ belongs to $C^{2,\eta}$ and solves the steady state equation $(L + \beta - \lambda_c)w = 0$. Then, also

$$(L + \beta - \lambda_c)h = 0. \quad (18)$$

Since (18) holds, X^H is a $(L + a \frac{\nabla h}{h} \cdot \nabla, 0, \alpha h e^{-\lambda_c t}; D)$ -superdiffusion by Lemma 1(b). From now on $\widetilde{\mathbf{P}}$ will denote the probability corresponding to X^H . By recalling the definition of $\mathfrak{S}$ and defining

$$\nu := h\mu, \quad g := f/h, \quad q := hr,$$

one can reformulate (A.2)(c) as follows:

$$(A^*.2)(c) \lim_{t \rightarrow \infty} \langle s_t \cdot \mathfrak{S}_t(g)(x), \nu(dx) \rangle = \langle g, q(dy) \rangle \cdot \|\nu\|, \quad g \in C_c^+(D), \quad \nu \in \mathcal{M}_f.$$

Similarly, (A.3)(d)-(A.3)(e) become

$$(A^*.3)(d) \lim_{t \rightarrow \infty} \widetilde{P}^\nu(\langle W_t, B_{z_t}^c(0) \rangle > 0) = 0, \text{ and}$$

$$(A^*.3)(e) \lim_{t \rightarrow \infty} \sup_{|x| \leq z_t} |s_{\widehat{z}_t} \cdot \mathfrak{S}_{\widehat{z}_t}(g)(x) - \langle g, q(dy) \rangle| = 0, \quad g \in C_c^+(D).$$

Finally, the theorem itself transforms into the following statement: if $g \in C_c^+(D) \not\equiv 0$ and $\nu \in \mathcal{M}_c(D) \not\equiv 0$, then in probability,

$$\lim_{t \rightarrow \infty} \frac{\langle X_t^H, g \rangle}{\widetilde{\mathbf{E}}^\nu \langle X_t^H, g \rangle} = \frac{\overline{W}_\infty}{\|\nu\|},$$

or, equivalently, in probability,

$$\lim_{t \rightarrow \infty} s_t \langle W_t, g \rangle = \langle g, q(dy) \rangle \overline{W}_\infty. \quad (19)$$

(In the equivalence we used that $\mathfrak{S}$ is the *expectation* semigroup along with (A*.2)(c).)

In order to show (19), the main idea is to use the comparison with the deterministic flow as in [8], nevertheless, there is an essential difference. In [8] we argued that by considering some large time $t+T$ (where both t and T are large), the changes of $\overline{X}^H$ are negligible after time t , while the remaining time T is still long enough to distribute the produced mass according to the ergodic flow given by the H -transformed migration. We then let first T and then t go to infinity.

Reading carefully the proof in [8] one can see that this method relied heavily on the ergodicity of the flow and would break down here. Hence, instead of letting first T and then t go to infinity, we now define

$$T := T_t = \widehat{z}_t.$$

Similarly to [8], the strategy is to

- (a) show that the total mass more or less stabilizes by time T_t ,
- (b) identify the limit of the *scaled flow* (starting from the state of the process at T_t) at time $t+T_t$
- (c) show that it agrees with the scaling limit of the measure-valued process itself.

Of course, (a) is simple: being a supermartingale, the *total mass converges*:

$$\lim_{t \rightarrow \infty} \|W_t\| = \overline{W}_\infty, \quad \tilde{\mathbf{P}}^\nu - a.s. \quad (20)$$

Unlike in [8], however, we do not know *a priori*, that the limit is non-zero, and moreover, one cannot proceed further without exploiting what is known about the radial speed of the process. Therefore we continue as follows. Let $\{Z_{W_t}(s)\}_{s \geq 0}$ denote the deterministic flow starting from the (random) measure W_t . Since given W_t ,

$$\langle Z_{W_t}(\widehat{z}_t), g \rangle = \langle \mathfrak{S}_{\widehat{z}_t}(g)(x), W_t(dx) \rangle,$$

one has

$$\tilde{\mathbf{P}}^\nu (|s_{\widehat{z}_t} \langle Z_{W_t}(\widehat{z}_t), g \rangle - \|W_t\| \langle g, q(dy) \rangle| > \epsilon) \leq A_t + B_t, \quad (21)$$

where

$$A_t := \tilde{\mathbf{P}}^\nu \left[\|W_t\| \cdot \sup_{|x| \leq z_t} |s_{\hat{z}_t}(\mathfrak{S}_{\hat{z}_t}(g))(x) - \langle g, q(dy) \rangle| > \epsilon \right],$$

$$B_t := \tilde{\mathbf{P}}^\nu [W_t(B_{z_t}^c(0)) > 0].$$

By (A*.3)(d-e), one has $\lim_{t \rightarrow \infty} A_t = \lim_{t \rightarrow \infty} B_t = 0$. Hence,

$$\lim_{t \rightarrow \infty} \tilde{\mathbf{P}}^\nu (|s_{\hat{z}_t} \langle Z_{W_t}(\hat{z}_t), g \rangle - \|W_t\| \langle g, q(dy) \rangle| > \epsilon) = 0. \quad (22)$$

From this, along with (A.3)(c) and (20), one obtains the *scaling limit of the flow*:

$$\lim_{t \rightarrow \infty} \tilde{\mathbf{P}}^\nu (|s_{t+\hat{z}_t} \langle Z_{W_t}(\hat{z}_t), g \rangle - \overline{W}_\infty \langle g, q(dy) \rangle| > \epsilon) = 0. \quad (23)$$

Our goal is therefore to show that *the scaling limit of the flow agrees with the scaling limit of the measure-valued process*. To achieve this, recall (A.2)(b). A computation using Chebysev's inequality and the supermartingale property (essentially the same computation as the one giving formula (28) in [8]) yields:

$$\begin{aligned} \tilde{\mathbf{P}}^\nu (|s_{t+\hat{z}_t} \langle Z_{W_t}(\hat{z}_t), g \rangle - s_{t+\hat{z}_t} \langle W_{t+\hat{z}_t}, g \rangle| > \epsilon) &\leq C \frac{\tilde{\mathbf{E}}^\nu \|W_t\|}{\epsilon^2 \lambda_c e^{\lambda_c t}} \cdot s_{t+\hat{z}_t}^2 \\ &\leq C \frac{\|\nu\|}{\epsilon^2 \lambda_c e^{\lambda_c t}} \cdot s_{t+\hat{z}_t}^2, \end{aligned}$$

where $C = C(\|g\|, \|\alpha h\|) := 18 \|\alpha h\| \cdot \|g\|^2$. Recall the abbreviation $T := \hat{z}_t$. If we show now that

$$\lim_{t \rightarrow \infty} e^{-\lambda_c t} s_{t+T}^2 = 0,$$

then we are done, since this implies

$$\lim_{t \rightarrow \infty} \tilde{\mathbf{P}}^\nu (|s_{t+T} \langle Z_{W_t}(T), g \rangle - s_{t+T} \langle W_{t+T}, g \rangle| > \epsilon) = 0.$$

To estimate $e^{-\lambda_c t} s_{t+T}^2$, observe that its logarithm can be estimated by the first condition in (A.2)(a):

$$\log(e^{-\lambda_c t} s_{t+T}^2) = -\lambda_c t + 2 \log s_{t+T} \leq -\lambda_c t + 2k \log(t+T),$$

with some $k > 0$ and, by (A.3)(b),

$$\lim_{t \rightarrow \infty} [-\lambda_c t + 2k \log(t+T)] = \lim_{t \rightarrow \infty} t [-\lambda_c + 2k \log(t+T)/t] = -\infty.$$

This completes the proof. $\square$

4.2 Proof of Theorem2

In [6] the following was shown.

Proposition 2 ([6], Theorem 3) *Assume that the diffusion process corresponding to L on D is not conservative and that*

$$\sup_{x \in D} \alpha(x) < \infty \text{ and } \inf_{x \in D} \beta(x) > -\infty. \quad (24)$$

Then the compact support property does not hold.

(In [6] this was stated for $D = \mathbb{R}^d$, but the proof goes through for general D too.)

Using this result, Theorem 2 follows easily by applying an h -transform. Indeed, let us suppose that X possesses the compact support property but the diffusion corresponding to L_0^h is not conservative. Since the support of the superprocess (and thus the compact support property too) is invariant under h -transforms, therefore X^h possesses the compact support property too. On the other hand, since X^h is the $(L_0^h, \lambda_c, \alpha h; D)$ -superprocess, L_0^h is not conservative, and αh is bounded from above, Proposition 2 implies that the compact support property does not hold; contradiction. $\square$

4.3 Proof of Theorem3

Recall that in our notation x is the d -dimensional vector $(x_1, x_2, \dots, x_d)$ and let $h(x) := \exp(cx_1)$. A straightforward computation reveals that X^h is the superprocess corresponding to the operator

$$\left[\frac{1}{2} \Delta + c \frac{\partial}{\partial x_1} + \left(\beta + \frac{c^2}{2} \right) \right] u - \alpha h u^2,$$

and starting at $\nu := h\mu$. Therefore, X^h in a coordinate system

$$x'_1 = x_1 - ct, \quad x'_2 = x_2, \dots, \quad x'_d = x_d$$

is equal in distribution to the super-Brownian motion corresponding to the operator

$$\left[\frac{1}{2} \Delta + \left(\beta + \frac{c^2}{2} \right) \right] u - \alpha h u^2.$$

(This can be derived easily for example from the log-Laplace equation (2).) By Theorem 1, this latter one obeys the Law of Large Numbers because $\alpha(x) \exp(cx_1)$ is bounded and $c < \sqrt{2\beta}$ guarantees that $\lambda_c > 0$. (Of course

the limiting random variable depends on c .) For X^h this means that if $f \in C_c^+(D) \not\equiv 0$ and $\mu \in \mathcal{M}_c(D) \not\equiv 0$, then in probability,

$$\lim_{t \rightarrow \infty} \frac{\langle X_t^h, f^{(ct)} \rangle}{\tilde{\mathbf{E}}^\nu \langle X_t^h, f^{(ct)} \rangle} = \frac{\overline{W}_\infty}{\|\nu\|},$$

where

$$f^{(ct)}(x) := f(x_1 + ct, x_2, \dots, x_d),$$

and $\overline{W}_\infty = \overline{W}_{\infty, c}$. Denoting $g := hf$, we obtain that if $g \in C_c^+(D) \not\equiv 0$ and $\mu \in \mathcal{M}_c(D) \not\equiv 0$, then in probability,

$$\lim_{t \rightarrow \infty} \frac{\langle X_t, g^{(ct)} \rangle}{\mathbf{E}^\mu \langle X_t, g^{(ct)} \rangle} = \frac{\overline{W}_\infty}{\langle h, \mu \rangle},$$

where $g^{(ct)}(x) := g(x_1 + ct, x_2, \dots, x_d)$. □

5 Appendix

In this section we provide some background material regarding various probabilistic and analytic concepts. These can be found more completely in [5, 8, 10].

5.1 The particle picture for the superprocess

In the introduction we defined the $(L, \beta, \alpha; D)$ -superprocess X analytically. In fact, X also arises as the short life time and high density diffusion limit of a *branching particle system*, which can be described as follows: in the n^{th} approximation step each particle has mass $1/n$ and lives a random time which is exponential with mean $1/n$. While a particle is alive, its motion is described by a diffusion process corresponding to the operator L . At the end of its life, the particle dies and is replaced by a random number of particles situated at the parent particle's final position. The distribution law of the number of descendants is spatially varying such that the mean number of descendants is $1 + \frac{\beta(x)}{n}$, while the variance is assumed to be $2\alpha(x)$. All these mechanisms are independent of each other.

See Appendix A in [5] for a precise statement on the particle approximation.

5.2 The generalized principal eigenvalue

Let Y be the diffusion process on D corresponding to L , and denote by $\mathbb{P}^x$ the law of Y starting at $x \in D$. Then from a probabilistic point of view, the generalized principal eigenvalue can be equivalently expressed as

$$\lambda_c = \sup_{\{A: A \subset \subset D, \partial A \text{ is } C^{2,\eta}\}} \lim_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}^x \left[\exp \left(\int_0^t \beta(Y_s) ds \right); \tau^A > t \right], \quad (25)$$

for any $x \in D$, where $\tau^A := \inf\{t \geq 0 : Y_t \notin A\}$, and the $C^{2,\eta}$ -boundary is defined with the help of $C^{2,\eta}$ -maps in the usual way. (See Section 4.4 in [10] on the subject). Hence, since β is bounded from above, $\lambda_c < \infty$; and it is known from standard theory that for any $\lambda \geq \lambda_c$, there exists a function $0 < f \in C^{2,\eta}(D)$ such that $(L + \beta)f = \lambda f$ on D . (See Section 4.3 in [10].)

5.3 Uniform integrability and conservativeness

Just like in [8], we show that

$$\lim_{t \rightarrow \infty} \widetilde{\text{Var}}^{h\mu}(\overline{X}^H) = \int_0^\infty ds e^{-2\lambda_c s} \langle \mu, \mathcal{S}_s[\alpha h^2] \rangle < \infty. \quad (26)$$

We will also see that if the diffusion process $\widehat{Y}$ corresponding to L_0^h on D is conservative, then $\overline{X}^H$ is a uniformly integrable $\widetilde{\mathbf{P}}^{h\mu}$ -martingale, whereas in general, $\overline{X}^H$ is a (non-negative) $\widetilde{\mathbf{P}}^{h\mu}$ -supermartingale. Clearly, only in the latter case it becomes an issue whether the limit is non-zero.

Indeed, if $\widehat{Y}$ is conservative, then consider the class

$$\mathfrak{C}(D) := \{f \in C^2(D) : \exists \mathcal{U} \subset D \text{ bounded s.t. } \overline{\mathcal{U}} \subset D; f = \text{const on } D \setminus \mathcal{U}\}.$$

By Lemma 1(b) along with Theorem A2 in [5], we have that for all $f \in \mathfrak{C}(D)$,

$$d\langle X_t^H, f \rangle = \langle X_t^H, L_0^h f \rangle dt + dM_t(f), \quad (27)$$

where $\{M_t(f)\}_{t \geq 0}$ is a square-integrable $\widetilde{\mathbf{P}}^{h\mu}$ -martingale, and its quadratic variation $\langle M(f) \rangle$ is given by

$$\langle M(f) \rangle_t = \int_0^t ds e^{-\lambda_c s} \langle X_s^H, \alpha h f^2 \rangle, \quad t \geq 0. \quad (28)$$

(The point is that one can take the function class $\mathfrak{C}(D)$ instead of just $C_c^2(D)$ when $\widehat{Y}$ is conservative.)

Applying (27) to the function $f \equiv 1$, it follows that $\overline{X}^H$ is a $\tilde{\mathbf{P}}^{h\mu}$ -martingale. Furthermore, by (28),

$$\tilde{\mathbf{E}}^{h\mu} [\langle X_t^H, 1 \rangle^2] = \langle \mu, h \rangle^2 + \int_0^t ds e^{-\lambda_c s} \langle h\mu, \mathfrak{S}_s[\alpha h] \rangle. \quad (29)$$

That is

$$\widetilde{\text{Var}}^{h\mu}(\overline{X}_t^H) = \int_0^t ds e^{-\lambda_c s} \langle h\mu, \mathfrak{S}_s[\alpha h] \rangle = \int_0^t ds e^{-2\lambda_c s} \langle \mu, S_s[\alpha h^2] \rangle. \quad (30)$$

Letting $t \rightarrow \infty$ we obtain (26). Replacing t by ∞ in the first of the integrals in (30), we have from (17) and from our assumptions that

$$\widetilde{\text{Var}}^{h\mu}(\overline{X}_t^H) \leq \int_0^\infty ds e^{-\lambda_c s} \langle h\mu, \mathfrak{S}_s[\alpha h] \rangle \leq \lambda_c^{-1} \|\alpha h\|_\infty \langle \mu, h \rangle < \infty.$$

Hence, by (29), $\sup_{t \geq 0} \tilde{\mathbf{E}}^{h\mu}(\overline{X}_t^H)^2 < \infty$, and consequently $\overline{X}^H$ is uniformly integrable.

Let $\mathfrak{D} := D \cup \{\Delta\}$ be the *one-point compactification* of D (when the underlying diffusion process $\hat{Y}$ is non-conservative on D , Δ is the *cemetery state* for $\hat{Y}$). Relaxing the assumption on the conservativeness of $\hat{Y}$, the argument in [5], pp. 726–727 shows that, although one can *not* work directly with the function class $\mathfrak{C}(D)$ (only with its subclass $C_c^2(D)$), extending X^H with $\tilde{\mathbf{P}}$ appropriately onto $\mathfrak{D}$ with $\mathfrak{P}$ makes $\overline{X}^H$ a $\mathfrak{P}^{h\mu}$ -martingale. Now, since the mass on the cemetery state Δ is nondecreasing in time, therefore $\overline{X}^H$ is less than this martingale by a non-decreasing process, that is, it is a $\tilde{\mathbf{P}}^{h\mu}$ -*supermartingale*. (In the non-conservative case, intuitively, mass is ‘lost’ at the Euclidean boundary of D or at infinity.)

Acknowledgement. The author owes thanks to J. Biggins for a helpful discussion regarding the proofs in [1] and [12].

References

- [1] **Biggins, J. D. (1992)** Uniform convergence of martingales in the branching random walk. *Ann. Probab.* **20(1)**, 137–151.
- [2] **Dawson, D. A. (1993)** Measure-valued Markov processes. In P.L. Hennequin, editor, *École d’été de probabilités de Saint Flour XXI–1991*, volume 1541 of *Lecture Notes in Mathematics*, pages 1–260. Springer-Verlag, Berlin.

- [3] **Dynkin, E. B. (1991)** Branching particle systems and superprocesses, *Ann. Probab.*, **19(3)**, 1157–1194.
- [4] **Dynkin, E. B. (2002)** Diffusions, superdiffusions and partial differential equations, *American Mathematical Society Colloquium Publications*, **50**. American Mathematical Society, Providence, RI.
- [5] **Engländer, J. and Pinsky, R. G. (1999)** On the construction and support properties of measure-valued diffusions on $D \subset \mathbb{R}^d$ with spatially dependent branching, *Ann. Probab.*, **27(2)**, 684–730.
- [6] **Engländer, J. and Pinsky, R. G. (2006)** The compact support property for measure-valued processes, *Ann. Inst. H. Poincaré Probab. Statist.* **42(5)** 535–552.
- [7] **Engländer, J. and Turaev, D. (2002)** A scaling limit theorem for a class of superdiffusions, *Ann. Probab.* **30(2)**, 683–722.
- [8] **Engländer, J. and Winter, A. (2006)** Law of large numbers for a class of superdiffusions, *Ann. Inst. H. Poincaré Probab. Statist.* **42(2)**, 171–185.
- [9] **Overbeck, L. (1994)** Some aspects of the Martin boundary of measure-valued diffusions. In *Measure-valued processes, stochastic partial differential equations, and interacting systems*, 179–186, AMS, Providence, RI.
- [10] **Pinsky, R. G. (1995)** Positive harmonic functions and diffusion. Cambridge University Press.
- [11] **Pinsky, R. G. (1995)** On the large time growth rate of the support of supercritical super-Brownian motion. *Ann. Probab.*, **23**, 1748–1754.
- [12] **Watanabe, S. (1967)** Limit Theorems for a Class of Branching Processes. In *Markov Processes and Potential Theory* (ed. J. Choquet), 205–232. Wiley, New York.